

Please find below the correct answers to the 2018 Academy Seminars individual learning assessment questionnaire:

1. To calculate the loading dose (LD) we need to know:

- a) Rate of absorption (K_a)
- b) Rate of elimination (K_{el})
- c) Volume of distribution (V_d)
- d) Area Under the Curve (AUC)

2. To calculate the maintenance dose (MD) we need to know:

- a) Time to reach the steady state (T_{ss})
- b) Volume of distribution (V_d)
- c) Clearance (CL)
- d) Rate of absorption (K_a)

3. If half life is 8 hours how long it takes to reach the steady state?

- a) 16 hours
- b) 24 hours
- c) 40 hours
- d) 48 hours

4. Indicate one of the main advantages of chromatographic methods over immunoassays.

- Low sample volume
- Less time of analysis
- Less cost
- More sensitivity
- Minimum sample preparation

5. Which of the following answers is false?

- Linear regression methods allow predictions of future concentrations with high reliability
- Bayesian methods should be used after validating the PK population models
- Non-linear regression methods need computers for their use

6. In pharmacotherapy, what is the inappropriate use of pharmacogenomics?

- Select the start dose for some drugs
- Select the appropriate drug in some type of cancer
- Useful to adjust dose for some drugs
- Useful to control toxicity

7. Blood/plasma sampling for TDM should always be done as a peak and trough level.

YES/NO

8. The best method of calculation of pharmacokinetic parameters is by using:

- o 1. The least squares method according to Sawchuck and Zaske
- o 2. The MAP Bayesian approach
- o 3. Standard two stage

9. The therapeutic range of a drug is a fixed value particular to that specific drug

YES/NO

10. What is the most important PK parameter for the first dose of an antibiotic?

- Clearance
- Volume of distribution

11. How long is the Post Antibiotic Effect or Post MIC Effect of an aminoglycoside?

- 2 hours
- 7 hours

12. A flucloxacillin serum concentration of 40 mg/L is enough to treat an infection with a micro-organism that has a MIC value of 2 mg/L

YES/NO (flucloxacillin is 95% protein bound and thus the free concentration is 2 mg/L. Optimal effect is seen at 4x the MIC = 8 mg/L)

13. After initialization of antiepileptic drug treatment blood concentrations should be measured after achievement of steady state which is attained by about 90% after 4 elimination half lives?

YES/NO

14. An indication to request TDM for typical antipsychotic drugs is weight gain due to supratherapeutic drug concentrations.

YES/NO

15. Smoking may affect the clearance of drugs by inducing drug metabolizing enzymes, especially CYP1A2. After smoking cessation the induced status "normalizes" within

- a) three days
- b) three weeks
- c) three months

16. Will TDM software tools improve the effectiveness of a TDM service?

YES/NO

17. Do modern TDM tools require steady state conditions for valid analysis and forecasts?

YES/NO

18. What is the benefit of Bayesian dose calculation?

- a. Allows a reduced number of samples
- b. The most likely result is obtained
- c. The results are less biased

19. The AUC is the best predictor of efficacy and/or toxicity for immunosuppressive and oncolytic agents

TRUE/FALSE

20. Please choose the correct statement(s):

Statement 1: Immunosuppressive agents can cause malignancies

Statement 2: Immunosuppressive agents can be used as oncolytic agents

a) Statements 1 and 2 are true

b) Statements 1 and 2 are false

c) 1 is true and 2 is false

d) 2 is true and 1 is false

21. PGx (pharmacogenetics) is related to clinical outcomes in transplantation.

TRUE/FALSE

Last update: 22 February 2019
